

## INDEX

*Note: In this index, the most detailed entries are under the scientific name (Latin binomial); vernacular names list page numbers only.*

### Abundance

CPUE and, 232–233

relative, 72

Active vs. passive (fixed) fishing gear, 267, 270, 306

Adelphophagy, 137

Aeterna Laboratories, 330

### Age

absolute vs. periodicity of ring formation, 116

defined, 99

at first reproduction, 145

at maturity, 143–145

Age bias plot, 113

Age-dependent mortality estimation methods, 173–174

Age determination, 105–115

ageing protocols, 106–107

back-calculation methods, 113–115

background and principles, 99–100

deviations and, 106–107

field sampling and storage, 101–102

reader agreement, precision and bias, 111–113

sampling coverage, 122–123

specimen collection and preparation, 100–105

staining protocols, 108–111

verification, 115–119

Web-based resources, 123

Age distribution, 191–192

Age-independent mortality estimation, 168–173

Age-structured matrix model (Leslie matrix), 195–197

Akaike's Information Criterion, 66

Allozymes, 83–84

*Alopias pelagicus* (pelagic thresher shark), 136

*Alopias superciliosus* (bigeye thresher shark), 136  
 Alopiidae (thresher sharks), 48  
 Amplified Fragment Length Polymorphism (AFLP), 88–89  
*Anacanthobatus* (leg skates), 51  
 Anal fin, family key, 38, 41  
 Analysis of molecular variation (AMOVA), 81–82  
 Angel sharks (Squatiniiformes), 46–47, 289  
     fecundity, 154  
     Pacific (*Squatina californica*), 84, 99, 141–142  
 Annuli (rings), 105, 107  
 Aplacental viviparity, 136  
 Aplacental yolk sac, 138  
*Aptychotrema rostrata* (Eastern Australian shovelnose ray), 152  
 Arches  
     haemal (transverse processes), 102  
     neural, 102, 110–111  
 Archival tags, 74–75  
     pop-up, 74–75  
 Area restrictions, 307–308  
 ARIMA (auto-regressive integrated moving average) models, 277  
 Arlequin software, 82  
 Artisanal fishing, 247  
 Assumption violation, 179  
 Atlantic sharpnose shark (*Rhizoprionodon terraenovae*), 85, 87, 151, 257  
 Atlantic stingray (*Dasyatis sabina*), 137  
 Atlantic yellowfin tuna (*Thunus albacares*), 12  
 At-sea vs. shoreside sampling, 258–262  
 Australian Cooperative Game-Fish Tagging Program, 63  
 Australian sharpnose shark (*Rhizoprionodon taylori*), 146–147, 153, 169, 175  
 Auto-regressive integrated moving average (ARIMA) models, 277  
 Availability, 292  
 Average percent error (APE) technique, 111–112  
  
 Back-calculation  
     in age determination, 113–115  
     Dahl-Lea direct proportions method, 114

linear-modified Dahl-Lea method, 114  
 quadratic-modified Dahl-Lea method, 114–115  
 size-at-birth-modified Fraser-Lea equation, 115  
 Bag limits, 304  
 Banding, centrum, 99–100  
 Barbeled houndsharks (*Leptochariidae*), 49  
 Basking shark (*Cetorhinus maximus*), 1, 48  
*Bathyraja brachyurops* (broadnose skate), 101  
 Batoids, 23, 46–47, 289  
     consumptive uses, 326  
     reproductive strategy, 135  
     vs. sawsharks, 50  
 Bayesian estimation, 230–231  
 Beverton and Holt recruitment model, 219  
 Bias, gear, 273–274  
 Bigeye thresher shark (*Alopias superciliosus*), 136  
 Biological objectives, 15–16  
 Biological productivity, 290–292, 295–296  
 Biological reference points (BRPs), 13  
 Biomass dynamics models, 295  
 Birthmark ring, 107  
 Blacktip sawtail catshark (*Galeus sauteri*), 145  
 Blacktip shark (*Carcharhinus limbatus*), 85, 90, 142, 153, 180  
 Blacktip shark (*Carcharhinus tilstoni*), 179  
 Blind sharks (*Brachaeluridae*), 47–48  
 Bluefin tuna, southern (*Thunnus maccoyii*), 74  
 Blue shark (*Prionace glauca*), 66, 87, 141, 151, 249, 326  
 Bluntnose stingray (*Dasyatis say*), 153  
 Body measurements, 255–257  
     family key, 42  
 Bomb carbon dating, 116, 118  
 Bonnethead shark (*Sphyrna tiburo*), 169, 175  
*Brachaeluridae* (blind sharks), 47–48  
 Bramble sharks (echinorhinids), 46  
 Brander's equilibrium mortality estimation, 170–173  
 Broadnose skate (*Bathyraja brachyurops*), 101

Brody equation, 216–217

Bullhead sharks (Heterodontiformes), 47

Butterfly rays (Gymnuridae), 51–52

Buy-back schemes, 302

Bycatch, 244, 307

Bycatch fisheries, 289–290

Bycatch Reduction Devices (BRDs), 247

  

CAGEAN model, 220, 224–228

- Doubleday method, 228
- Fournier and Archibald method, 228
- Paloheimo method, 225–228

Calcein validation, 118

Calcification, 99–100

- of neural arches, 110–111

Callorhynchidae (elephant fishes), 52

Capture-recapture studies, 176, see also Tag-recovery studies

*Carcharias taurus* (sand tiger shark), 2

Carcharhinidae (requiem sharks), 49–50

Carcharhiniformes (ground sharks), 49–50, 135, 136

*Carcharhinus isodon* (fine-tooth shark)

- fecundity, 154
- reproductive cycle, 151
- staining methods, 109, 110

*Carcharhinus limbatus* (blacktip shark)

- male maturity, 142
- mortality estimation, 180
- reproductive strategy, 153

*Carcharhinus obscurus* (dusky shark), 1, 2, 249

*Carcharhinus plumbeus* (sandbar shark), 65, 85, 142

- matrix modeling, 198–199
- mortality estimation, 174
- sexual segregation, 257

*Carcharhinus signatus* (night shark), 2

*Carcharhinus sorrah* (spot-tail shark), staining method, 111

*Carcharhinus tilstoni* (blacktip shark), 85, 90, 179

*Carcharias taurus* (grey nurse shark), sanctuaries, 309

*Carcharias taurus* (sand tiger shark)

- oxytetracycline validation, 118
- reproductive interval, 152
- reproductive strategy, 137
- survival time, 249

*Carcharodon carcharias* (white shark), 87, 90

Carpet sharks (Orectolobiformes), 47–48

Cartilage extracts, 330

Cartilage pills, 329–330

Cataloguing, 31

Catchability, 292, 293

Catch-and-release fishing, 332–333

Catch-at-age analysis, see CAGEAN model

Catch curves, in mortality estimation, 174–176

Catch estimates, 243–244, see also CPUE

- bycatch, 244
- disposition estimates, 244

Catch Per Unit Effort, see CPUE

Catch quotas, 304–305

Catch rate

- by artisanal fishing, 247
- in depths of 25–50 M, 246–247
- sharks vs. other species, 246

Catch time series, 251

Catshark, 47–48, 49

- blacktip sawtail (*Galeus sauteri*), 145
- chain (*Scyliorhinus retifer*), 154
- false (*Pseudotriakis microdon*), 49, 137
- finback (Proscylliidae), 49
- small-spotted (*Scyliorhinus canicula*), 145, 154

Caudal fin, family key, 34, 36–37, 39, 40, 41, 42

Caudal thorns, in age and growth studies, 101

Centre for Environment, Fisheries and Aquaculture Science (CEFAS), 74

Centrophorids (gulper sharks), 24, 46

*Centroscyllium fabricii* (deep sea black dogfish), 145

Centrum banding, 99–100, 105

Centrum edge analysis, 116–117

*Cetorhinus maximus* (basking shark), 1, 48, 168–173

Chain catshark (*Scyliorhinus retifer*), 154

Chemical tagging, 116

Chen and Watanabe mortality estimation method, 173–174

Chimaeriformes (chimaeras), 23, 25, 52, 289

*Chirrhigaleus*, 46

Chondrichthyans, 23–24

Claspers, 138–139, 157

- length at maturity, 143–144
- maturity and, 141

Closed periods, 302

Codes for species identification, 255

Coffin rays (Hypnidae), 50

Cohort analysis

- CAGEAN, 220, 224–228
- Virtual Population Analysis, 180, 220–224

Common skate (*Dipturus batis*), 63

Conceptual models, 13–15

Conservation objectives, 15–16

Contrast, in data, 232–233

Convention for International Trade in Endangered Species (CITES), 3, 298

Cownose rays (Rhinopteridae), 51–52

CPUE, 211

- abundance and, 232–233
- in CAGEAN, 226
- data collection, 245–248, see also Fishing gear
- definition, 244–245
- in delay-difference model, 218
- in surplus-production model, 211–212

Crocodile sharks (Pseudocarchariidae), 48

Crystal violet protocol, 109–110

Customary Marine Tenure (CMT), 300–301

Dahl-Lea direct proportions method, 114

*Dalatias*, 46

Dart tag, 63

Dasyatidae (stingrays), 51–52

*Dasyatis centroura*, 152

*Dasyatis marmorata*, 152

*Dasyatis sabina* (Atlantic stingray), 137

*Dasyatis say* (bluntnose stingray), 153

Data, contrast in, 232–233

Data auditing software, 280

Database creation, 278–280

Database software, 279

Data collection, 63–73, see also Tag recovery studies  
for species-specific sampling, 253–254

Data quality, 231–233

Data storage (archival) tags, 74–75

Decommissioning schemes, 302

Deep sea black dogfish (*Centroscyllium fabricii*), 145

Delay-difference (Deriso) model, 215–220  
advantages and disadvantages, 219–220  
examples, 220

Demographic models, 187–203  
background and principles, 189–190  
conclusions and advice, 200–201  
life tables, 190–195, see also Life tables  
matrix models, 195–200, see also Matrix models

Devil rays (Mobulidae), 51–52

*Dipturus batis* (common skate), 63

Disc, family key, 33, 36

Disposition estimates, 244

Dissection, 32

Diversity  
haplotype, 86  
nucleotide sequence, 86

Diving, recreational, 332

DNA

- mitochondrial, 83, 84–86
- nuclear, 83
- DNA microsatellites, 86–88
- Dockside sampling, 260–262
- Dogfish (Squaliformes), 46
  - deepsea black (*Centroscyllium fabricii*), 145
  - piked/spiny (*Squalus acanthias*), 106, 139, 140, 142, 148, 153, 155, 169, 257
  - smooth (*Mustelus canis*), 108, 109
- Dorsal fin, family key, 33, 38, 39, 44
- Dorsal fin spines
  - in age and growth studies, 101
  - preparation, 104–105
- Doubleday method, 228
- Dusky shark (*Carcharhinus obscurus*), 1, 2, 249
- Dusky smooth-hound (*Mustelus canis*), 138, 139, 145–146, 148, 151
- Dynamics, fishery, 11
  
- Eagle rays (Myliobatidae), 51–52
- Eastern Australian shovelnose ray (*Aptychotrema rostrata*), 152
- Echinorhinids (bramble sharks), 46
- Ecolabeling, 317
- Economic objectives, 16
- Ecosystem approach, 291–292
- Effort (quantitative) use rights, 302–304
- Elasmobranchs, 23
- Elasticity, 200
- Electric rays (Torpidiniformes), 50
- Elephant fishes (Callorhynchidae), 52
- Enclosure, fishing area, 310–311
- Encounterability, 292–293
- Endangered species, 315–316
- Environmental Analysis System (EASy), 280
- Equilibrium constraint, 207
- Equilibrium mortality estimation methods, 170–173
- Etmopterus granulosus* (southern lanternshark), 46, 154
- Exclusive Economic Zones, 299

Exploitation rates of mortality, 178–179

Exponential survival model, 214

Extinction, 2–3

Eyelid

family key, 44

nictating, 2y

Eyes, family key, 43–44

Fabens equation, 121

False catshark (*Pseudotriakis microdon*), 49, 137

Family key, 32–45

anal fin, 38, 41

body, 42

caudal fin, 34, 36–37, 39, 40, 41, 42

disc, 33, 36

dorsal fin, 33, 38, 39, 44

eyelid, 44

eyes, 43–44

flap, 37

gill openings, 33, 43

gill slits, 33, 36

head, 39

mouth, 38, 40, 43

nostril, 41

pelvic fin, 35, 39, 44

precaudal pit, 44

snout, 33, 35, 38, 39–40, 40, 45

spiracles, 44–45

tail, 42

thorns or fine denticles, 35–36

trunk, 39

Fecundity, 1, 153–155, 194, see also Reproductive biology

Field data collection, reproductive biology, 158–159

Field identification, 28–30

body measurements, 29

color, 28–29

- fin placement, 29
- teeth counts, 30
- vertebral counts, 29–30
- Finback catsharks (*Proscylliidae*), 49
- Finetooth shark (*Carcharhinus isodon*), 109, 110, 151, 154
- Fins, consumptive uses, 326–328
- First-growth rings, 107
- First-year survival, 194
- Fisheries
  - defined, 9
  - Individual Transferable Quota, 243–244
  - stock assessment examples of real, 236
- Fisheries management, see Management
- Fisheries observer programs, 258, 259
- Fisheries science, see also Management; Stock assessment
  - defined, 9
  - objective, 9–10
- Fishery-dependent sampling, 241–263, see also under CPUE; Sampling
- Fishery dynamics, 11
- Fishery-independent sampling, 265–274, see also under Sampling
- Fishing area closure, 308, 310–311
- Fishing gear, 288–289
  - gear bias, 273–274
  - gillnet, 245–246, 289
  - hook, 289
  - longline, 246–247
  - passive (fixed) vs. active, 267, 270, 306
  - purse seine, 248
  - regulation, 305–307
  - regulation of, 303, 305–307
  - survey design considerations, 273–276
  - trawl, 247–248, 289
- Fishing gear performance databases, 279
- Fishing licenses, 301, 302
- Fishing location, 250–251
  - catch time series, 251

- recording methods, 250–251
- Fishing mortality, 13, 192, 225–226, 249–250, 292–294
- Flap, family key, 37
- Fluorescein validation, 118
- Fournier and Archibald method, 228
- Fraser-Lea equation, size-at-birth-modified, 115
- Freshwater species, 290
- Frilled sharks, 45
- $F_{ST}$  statistic, 81
- Furgaleus macki* (whiskery shark)
  - reproductive interval, 152
  - sperm storage, 156
  
- Galeocerdo cuvier* (tiger shark), 66, 249
  - male maturity, 142
- Galeomorphi, 23
- Galeorhinus galeus* (school shark), 179, 180, 311, 315
- Galeorhinus galeus* (soupfin shark), 1
- Galeorhinus galeus* (tope shark), 66
  - fecundity, 154, 155
  - reproductive strategy, 153
  - sperm storage, 156
- Galeus sauteri* (blacktip sawtail catshark), 145
- GAMs (Generalized Additive Models), 278
- Gear bias, 273–274
- Gear performance databases, 279
- Gear selectivity, 69–70
- Genepop software, 82
- Generalized Additive Models (GAMs), 278
- Generalized linear models (GLMs), 277–278
- Genetics, 79–96, see also Molecular markers; Stock structure
- Giant guitarfish (*Rhynchobatus djidensis*), 328
- Gillnet fishing gear, 245–246, 273, 289
- Gill opening, family key, 33
- Gill slits, family key, 33, 36
- Ginglymostoma cirratum* (nurse shark), 249

Ginglymostomatidae (nurse sharks), 47, 87

Glossaries, see Terminology

Goblin sharks (Mitsukurinidae), 48

*Gollum attenuatus* (slender smooth-hounds), 137

Gompertz growth function, 119, 121–122

Gonadosomatic index (GSI), 146–147

GPS systems, 251

Ground sharks (Carcharhiniformes), 49–50

Growth

- in captive animals, 118
- vertebral, 99

Growth models, 119–122

- background and principles, 99–100
- field sampling and storage, 101–102
- Gompertz growth function, 119, 121–122
- sample size and, 119–120
- sampling coverage, 122–123
- verification, 115–119
- von Bertalanffy growth function, 119, 120–121
- Web-based resources, 123

Growth rates, 66–69

Guitarfish

- common (*Rhinobatos rhinobatos*), 151, 154
- giant (*Rhynchobatus djidensis*), 328
- sharkfin (Rhyncobatiformes), 51

Gulper sharks (centrophorids), 46

Gummy shark (*Mustelus antarcticus*), 1, 84, 89–90, 109, 137, 179, 180, 307

Gunderson–Dygert mortality estimation method, 169

Gunderson mortality estimation method, 169

Gymnuridae (butterfly rays), 51–52

Habitat utilization, 65

Haemal arches (transverse processes), in age and growth studies, 102

Hammerhead, 49, 87, 249

- scalloped (*Sphyrna lewini*), 155
- smalleye (*Sphyrna tudes*), 141–142, 153

Haplotype diversity, 86

Hardy-Weinberg equilibrium, 81–82

Head, family key, 39

Hemicyllidae (longtail carpet sharks), 47–48

Hemigaleidae (weasel sharks), 49

Hemiscyllids, 48

Heterodontiformes (bullhead or horn sharks), 47, 135

*Heterodontus portusjacksoni* (Port Jackson shark), 148

Hexanchiformes, 45

- reproductive strategy, 136

*Hexanchus griseus* (sixgill sharks), 111

Hexatrygonoidea (sixgill stingrays), 51–52

*Himantura*, 24

Histological examination, of reproductive tract, 147–150

Hoenig mortality estimation method, 169–170

Hook fishing gear, 273, 289

Hook size, 307

Houndshark, 49

- barbeled (Leptochariidae), 49

Hyperdepletion, 233

Hyperstability, 233

Hypnidae (coffin rays), 50

Identification

- field, 28–30
- laboratory, 30

Illumination, of sectioned centra, 105–106

Individual Transferable Quotas (ITQs), 304–305

Instantaneous rates of mortality, 178–179

Institutional frameworks, 298–299

Internal anchor tag, 61–62, 63

- body cavity type, 61–62
- button type, 62

International developments, management measures and, 298

International Plan of Action for the Conservation and Management of Sharks (IPOA-Sharks), 3, 298

Intrinsic rate of population growth, 209, 295

Island model of migration, 82–83

*Isurus oxyrinchus* (shortfin mako shark), 70

- bomb carbon dating, 118
- reproductive strategy, 136

IUCN Red List, 316

Japan, reporting classes, 252

Jensen mortality estimation method, 170

Jumbo Rototag, 62–63

Jurisdictional frameworks, 298–299

Kimura method, 120

Laboratory identification, 30

Lake Erie perch, Paloheimo method example, 226–228

*Lamna ditropis* (salmon shark), 87, 104

*Lamna nasus* (porbeagle shark), 1, 66, 87

- bomb carbon dating, 118
- mortality, 168–173
- mortality estimation, 180
- reproductive interval, 152
- reproductive strategy, 136

Lamniformes (mackerel sharks), 48–49, 139

Landings, carcasses, 249

Landings sampling, 248–249

Lanternshark, southern (*Etmopterus granulosus*), 154

Large ray (*Rhina ancylostoma*), 50

Latin binomial vs. vernacular names, 252

Leather, shark skin, 328–329

Leg skates (*Anacanthobatus*), 51

Lemon shark (*Negaprion brevirostris*), 65, 88, 105, 109, 195

Length/weight relationship, 66

Leopard shark (*Triakis semifasciata*), 111, 180, 194

Leptochariidae (barbeled houndsharks), 49

Leslie Matrix, 195–197

*Leucoraja erinacea* (little skate), 99

Licenses, 301  
     rescinding of, 302  
 Life tables, 190–195  
     Australian sharpnose shark (*Rhizoprionodon taylori*) example, 192–193  
     general approaches, 190–193  
     rebound potential, 194–195  
 Light, transmitted vs. reflected, 105–106  
 Lighting, of sectioned centra, 105–106  
 Likelihood ratio tests, 120  
 Limited entry use rights, 301–302  
 Linear models, generalized (GLMs), 233  
 Linear-modified Dahl-Lea method, 114  
 Linear regression, 229–230  
 Little skate (*Leucoraja erinacea*), 99  
 Liver, consumptive utilization, 330–331  
 Lobed stingaree (*Urolophus lobatus*), 156  
 Locally weighted regression scatter plot smoothing (LOWESS), 277  
 Logbooks, 260, 261  
 Longline fishing gear, 246–247  
 Longnose chimeras (Rhinochimaeridae), 52  
 Longtail carpet sharks (Hemiscylliidae), 47  
 LORAN systems, 251  
 LOWESS method, 277  
  
 Mackerel sharks (Lamnidae), 48–49  
 Magnuson-Stevens Fishery Conservation and Management Act (1996), 16  
 Mako, shortfin (*Isurus oxyrinchus*), 70, 85, 90–91, 118, 136  
 Management, 285–321  
     background and principles, 287–288  
     as balancing act, 17–18  
     basic concepts, 9–10  
     biological aspects, 288–298  
         fisheries impact, 288–290  
         fishing mortality, 292–294, see also Fishing mortality  
         risk evaluation, 294–297  
         species biology, 290–292

- ecosystem approach, 291–292
- frameworks for, 298–299
- objectives of, 15–18, see also Management objectives
- product certification and ecolabeling, 317
- technical, 305–315
  - area and time restrictions, 307–308
  - fishing area enclosure, 310–311
  - fishing gear regulation, 305–307
  - Marine Protected Areas (MPAs), 308–309
  - product form, 313–314
  - regional fisheries management, 311–313
  - size limits, 314–315
- use rights, 289–305
  - limited entry, 301–302
  - quantitative input (effort), 302–304
  - quantitative output (catch quotas), 304–305
  - territorial, 300–301
- Management objectives, 15–18
  - balance among, 17–18
  - biological and conservation, 15–16
  - economic, 16
  - in opposition, 15
  - recreational, 17
  - social, 17
- Management principles, 1–6
- Marginal Increment Ratio (MIR), 117
- Marine Protected Areas (MPAs), 308–309
- MARK software, 178
- Masked stingaree (*Trygonoptera personata*), 153
- Mating, 146, see also Reproductive cycle
- Matrix models
  - age-structured (Leslie matrix), 195–197
  - elasticities, 200
  - stage-based, 197–199
- Maturity, 140–145
  - age at, 143–145

- assessing, 140–141
  - female, 142–143
  - male, 141–142
- Maximum economic rent (MER), 16
- Maximum sustainable yield (MSY), 11–13, 194, 195
- Mean generation length, 191
- Megachasmidae (megamouth sharks), 48
- Mercury content, 326
- Mesh size, 307
- Microsatellites, 86–88
- Migration, island model of, 82–83
- Mitochondrial DNA, 83, 84–86
- Mitsukurinidae (goblin sharks), 48
- Mobulidae (devil rays), 51–52
- Model complexity, 13–15
- Molecular markers, 83–91
  - allozymes, 83–84
  - Amplified Fragment Length Polymorphism (AFLP), 88–89
  - case studies, 89–91
    - blacktip shark (*C. limbatus*), 90
    - gummy shark (*Mustelus antarcticus*), 89–90
    - shortfin mako, 90–91
    - white shark (*Carcharodon carcharias*), 90
  - microsatellites, 86–88
  - mitochondrial DNA, 83, 84–86
  - Random Amplified Polymorphic DNA (RAPD), 88
  - tissue collection, 89
- Mortality, 13, 72
  - exploitation rates of, 178–179
  - fishing, 192, 225–226, 249–250, 292–294
  - instantaneous rates of, 178–179
  - natural, 192, 295–296
  - post-capture, 292, 293
  - respiratory mode and, 249
  - types of, 167
- Mortality estimation, 165–185

- background and principles, 167–168
- conclusions and advice, 181–182
- direct methods, 174–181
- indirect methods, 168–174
  - age-dependent, 173–174
  - age-independent, 168–173
  - catch curves, 174–176
  - cohort analysis (Virtual Population Analysis), 180–181
  - tag recovery studies, 176–180
  - telemetry, 180
- Mortality rates
  - finite, 167
  - instantaneous, 167
- Mouth, family key, 38, 40, 43
- Movement, data collection, 70–72
- Multispecies modeling, 312
- Mustelus antarcticus* (gummy shark), 1, 84, 89–90
  - age determination, 109
  - mesh size regulation and, 307
  - mortality estimation, 180
  - reproductive strategy, 137
  - tag-recovery studies, 179
- Mustelus canis* (dusky smooth-hound)
  - age at first reproduction, 145
  - fecundity, 155
  - reproductive anatomy, 139
  - reproductive cycle, 145–146, 148, 151
  - reproductive strategy, 138
  - sperm storage, 156
- Mustelus canis* (smooth dogfish), age determination, 108, 109
- Mustelus griseus* (spotless smooth-hound), 138, 148
- Mustelus lenticulatus* (spotted estuary smooth-hound), 70, 137, 151
- Mustelus manazo*, 148
- Mustelus* spp., 24, 50
  - reproductive strategy, 137–138
- Myliobatidae (eagle rays), 51–52

Myliobatiformes (stingrays), 51–52

Myloibatoidei (stingrays), 51–52

Narcinidae (numbfishes), 50

Narkidae (sleeper rays), 50

National Marine Fisheries Service, see NMFS entries

National Plans of Action, 3, 298, 314

National Shark Assessment Reports, 3

Natural mortality, 192, 295–296

*Negaprion brevirostris* (lemon shark), 65, 88, 195

- age determination, 109
- banding patterns, 105

Neovastat, 330

Net reproductive rate, 191

Neural arches

- in age and growth studies, 102
- calcification, 110–111

Nictating eyelid, 27

Night shark (*Carcharhinus signatus*), 2

NMFS

- Candidate List for Threatened and Endangered Species, 2
- Cooperative Shark Tagging Program, 63
- Fishery Management Plan, 2

Nostril, family key, 41

Nucleotide sequence diversity, 86

Numbfishes (Narcinidae), 50

Nursery areas, delineation, 65

Nurse shark (*Carcharias taurus*), 309

Nurse shark (*Ginglymostoma cirratum*), 249

Nurse sharks (Ginglymostomatidae), 47, 87

Odontaspidae (sand tiger sharks), 48

Oophagy, 136–137

Orectolobidae (wobbegongs), 47–48

Orectolobiformes (carpet sharks), 47–48

- reproductive strategy, 135, 136

survival time, 249

ORI tag, 62–63

Osteichthyes, 23

Output control, 304–305

Oviparity, 135–136

Ovulation, 150

Oxynotids (prickly dogfishes), 46

Oxytetracycline validation method, 117–118

Pacific angel shark (*Squatina californica*), 84, 99, 141–142

Paloheimo method, 225–228

Parascylliidae (collared carpet sharks), 47–48

Parascylliids, 48

Passive (fixed) vs. active fishing gear, 267, 270, 306

Pauly mortality estimation method, 168

Pelagic thresher shark (*Alopias pelagicus*), 136

Pelvic fin, family key, 39, 44

Percent reader agreement, 112

Petersen disc tag, 63

Peterson and Wroblewski mortality estimation method, 173

Photography, 30–31

Piked/spiny dogfish (*Squalus acanthias*), 1, 106, 139, 140, 142, 148, 153, 155, 257

Placental viviparity, 137–138

Platyrrhinidae (thornback rays), 52

Platyrrhinoidei (stingrays), 51–52

Population growth

- finite rate of, 209
- intrinsic rate of, 295

Pop-up archival tags, 74–75

Porbeagle shark (*Lamna nasus*), 1, 66, 87, 118, 136, 152, 180

Port Jackson shark (*Heterodontus portusjacksoni*), 148

Post-capture mortality, 292, 293

Pre-birth rings, 107

Precaudal pit, family key, 44

Predator role, 2–3

Prickly dogfishes, 46

*Prionace glauca* (blue shark), 66, 87, 249, 326  
     reproductive cycle, 151  
     reproductive maturity, 141  
 Prior probability, 230  
 Pristiformes (sawfishes), 46, 50, 136  
 Pristiophoridae (sawsharks), 46, 50, 136  
 Probability  
     conditional, 230  
     prior, 230  
 Product certification, 317  
 Product form, 313–314  
 Profit maximization, 16  
 Proscylliidae (finback catsharks), 49  
 Pseudocarchariidae (crocodile sharks), 48  
 Pseudotriakidae (false catsharks), 49  
*Pseudotriakis microdon* (false catshark), 137  
 Purse seine fishing gear, 248  
  
 Quadratic-modified Dahl-Lea method, 114–115  
 Quality assurance/quality control, in surveys, 272  
 Quantitative input (effort) use rights, 302–304  
 Quantitative output (catch quotas) use rights, 304–305  
 Quantitative prediction, 13  
 Quotas, catch, 304–305  
  
*Raja clavata* (thornback ray), 74, 136  
     fecundity, 154  
 Rajidae, reproductive strategy, 135  
 Rajiformes (skates), 51  
 Rajimorphii, 23, see also Batoids  
 Random Amplified Polymorphic DNA (RAPD), 88  
 Rays, see also individual species  
     Eastern Australian shovelnose (*Aptychotrema rostrata*), 152  
     electric (Torpidiniformes), 50  
     taxonomic terminology, 25  
     thornback (*Raja clavata*), 136, 154

Reader agreement, in age determination, 111–112

Rebound potential, 194–195

Recreational objectives, 17

Recreational utilization, 332–333

Recruitment

- Beverton and Holt model, 219
- Ricker model, 218–219

Regional Fisheries Management Organizations (RFMOs), 299, 311–313

Relative abundance, 72

Relative marginal increment analysis (RMI), 117

Reproductive anatomy, 138–140

Reproductive biology, 133–164, see also Subtopics

- anatomy, 138–140
  - female, 139–140
  - male, 138–139
- fecundity, 153–155
- field data collection, 158–159
- maturity, 140–145
  - female, 142–143
  - male, 140–142
- reproductive cycle timing, 145–153
- reproductive strategies, 135–138
- resources, 156–158
- sperm storage, 155–156

Reproductive cycle, 145–153

- female, 150–153
  - examples, 152–153
  - gestation cycle and time of birth, 150–151
  - ovulation, 150
  - reproductive interval, 152
- histological examination, 147–150
- male, 146–147
- mating, 146

Reproductive data collection, 257–258

Reproductive strategies

- aplacental viviparity, 136

- adelphophagy, 137
- aplacental yolk sac, 138
- oophagy, 136–137
- placental analogues, 137
- oviparity, 135–136
- placental viviparity, 137–138
- Requiem sharks (Carcharhinidae), 49–50
- Rhina ancylostoma* (large ray), 50
- Rhincodontidae (whale sharks), 47
- Rhincodon typus* (whale shark), 309
- Rhinidae, 50
- Rhiniiformes (sharkrays), 50
- Rhinobatiformes, 136
- Rhinobatos rhinobatos* (common guitarfish)
  - fecundity, 154
  - reproductive cycle, 151
- Rhinochimaeridae (longnose chimeras), 52
- Rhinopteridae (cownose rays), 51–52
- Rhizoprionodon taylori* (Australian sharpnose shark)
  - Leslie matrix example, 196–197
  - life table example, 192–193
  - mortality estimation, 169, 175
  - reproductive cycle, 146–147
  - reproductive strategy, 153
- Rhizoprionodon terraenovae* (Atlantic sharpnose shark), 85, 87
  - reproductive cycle, 151
  - sexual segregation, 257
  - as VPA example, 222–224
- Rhynchobatidae, 50
- Rhynchobatus*, 50
- Rhynchobatus djidensis* (giant guitarfish), 328
- Rhynchobatiformes (wedgfishes, sharkfin guitarfishes), 51
- Ricker recruitment model, 218–219
- Rig (*Mustelus lenticulatus*), 70
- Rings (annuli), 105
  - birthmark, 107

first-growth, 107  
pre-birth, 107  
Risk assessment, 294–297  
Rototags, 62–63

Salmon shark (*Lamna ditropis*), 87, 104  
Sample size, von Bertalanffy growth function and, 121  
Sampling  
  for age and growth studies, 101–105  
  at-sea vs. shoreside, 258–262  
  cleaning, cutting, mounting vertebrae, 102–105  
  fishery-dependent, 241–263  
    catch estimates, 243–244  
    CPUE, 244–248  
  fishery-independent, 265–274, see also Surveys  
    basic theory, 269–273  
    fishing gear, 273–276  
    implementation, 273–276  
    statistical methods, 276–280  
    survey design, 271–273  
    utility of surveys, 267–269  
  fishing area, 250–251  
  fishing mortality, 249–250  
  landings, 248–249  
  Latin binomial vs. vernacular names, 252  
  sex, 257–258  
  size, 254–257  
  species identification, 252–254  
Sampling coverage, for age/growth studies, 122–123  
Sampling design, 64  
Sampling frame, 269–270  
Sandbar shark (*Carcharhinus plumbeus*), 65, 85, 142, 174, 198–199, 257  
Sand tiger shark (*Carcharias taurus*), 2, 48, 118, 137, 152, 249  
Sawfishes (Pristiiformes), 50  
Sawsharks (Pristophoriformes), 46, 50  
Scalloped hammerhead (*Sphyrna lewini*), 155

Schaefer surplus production model, 208–210

School shark (*Galeorhinus galeus*), 179, 180, 311, 315

Scyliorhinidae (catsharks), 47–48, 49

*Scyliorhinus canicula* (small-spotted catshark), 145, 154

*Scyliorhinus retifer* (chain catshark), 154

Sectioning, vertebral, 102–103

Selectivity, 293

Sevengill sharks, 45

Sex identification, 257

Sex segregation, 257

Sharkfin guitarfishes (Rhincobatiformes), 51

Sharkrays (Rhiniformes), 50

Sharks, taxonomic terminology, 24

Shark skin leather, 328–329

Sharpnose shark

- Atlantic (*Rhizoprionodon terraenovae*), 85, 87, 257
- Australian (*Rhizoprionodon taylori*), 169, 175, 192–193, 196

Shortfin mako shark (*Isurus oxyrinchus*), 70, 85, 90–91, 136

Shortnose chimaeras (Chimaeridae), 52

Shortspine sourdog (*Squalus mitsukurii*), 154

Silver nitrate protocol, 110–111

Sixgill shark (*Hexanchus griseus*), 45, 111

Size, 254–257

- fisheries targeting size classes, 255
- importance in sampling records, 254
- weights and morphological measurements, 255–257

Size-at-birth-modified Fraser-Lea equation, 115

Size limits, 255, 314–315

Skates (Rajiformes), 51

- broadnose (*Bathyraja brachyurops*), 101
- consumptive uses, 326
- little (*Leucoraja erinacea*), 99

Skinning, 328–329

Sleeper rays (Narkidae), 50

Sleeper sharks (somniosids), 46

Slender smooth-hound (*Gollum attenuatus*), 137

Smalleye hammerhead shark (*Sphyrna tudes*), 153

Small-spotted catshark (*Scyliorhinus canicula*), 145, 154

Smooth dogfish (*Mustelus canis*), 108, 109

Smooth-hound

- dusky (*Mustelus canis*), 138, 139, 145–146, 148, 151, 155, 156
- reproductive cycle, 148
- slender (*Gollum attenuatus*), 137
- spotless (*Mustelus griseus*), 138
- spotted estuary (*Mustelus lenticulatus*), 137, 151

Snout, family key, 33, 35, 38, 39–40, 40, 45

Social objectives, 17

Software

- data auditing, 280
- database, 279
- Excel POPTOOLS, 196
- molecular biology, 82
- for mortality estimation, 178

Somniosids (sleeper sharks), 46

Soupfin shark (*Galeorhinus galeus*), 1

Southern bluefin tuna (*Thunnus maccoyii*), 74

Southern lanternshark (*Etmopterus granulosus*), 154

Spatial and temporal distribution, 72

Species identification

- codes for, 255
- materials used for, 253
- problems in landings data, 248–249
- problems with, 252–253
- in sampling, 252–254

Specimen collection, 31

Specimen preservation, 31

Spermatocyst, 147

Sperm stages, of testis, 149

Sperm storage, 155–156

*Sphyrna lewini* (scalloped hammerhead), 155

*Sphyrna* spp. (hammerhead), 45, 87, 249

*Sphyrna tiburo* (bonnethead shark), 175

*Sphyrna tudes* (smalleye hammerhead)

- male maturity, 141–142
- reproductive strategy, 153

Spiny dogfish, see Piked/spiny dogfish; *Squalus acanthias*

Spiracles, family key, 44–45

SPlus program, 280

Spotless smooth-hound (*Mustelus griseus*), 138

Spot-tail shark (*Carcharhinus sorrah*), 111

Spotted estuary smooth-hound (*Mustelus lenticulatus*), 151

Squalamine, 331

Squalene, 331

Squaliformes (dogfishes), 46, 289

- reproductive strategy, 136

Squalomorphi, 23

*Squalus acanthias* (piked/spiny dogfish), 1

- age determination, 106
- fecundity, 155
- female maturity, 142
- mortality estimation, 169
- reproductive anatomy, 139, 140
- reproductive cycle, 148, 153
- sexual segregation, 257

*Squalus mitsukurii* (shortspine sourdog), 154

*Squalus* spp., 24, 46

*Squatina californica* (Pacific angel shark), 84

- growth and age, 99
- male maturity, 141–142

*Squatina oculata* (angel shark), 154

*Squatina squatina* (angel shark), 154

- fecundity, 154

Squatiniiformes (angel sharks), 46–47, 289

- reproductive strategy, 136

Stable age distribution, 191–192

Stage-based matrix models, 197–199

Staining methods, 108–111

- crystal violet protocol, 109–110
- silver nitrate protocol, 110–111
- Statistical estimation, 276–280
  - database creation, 278–280
  - influence of covariates, 278
  - smoothing procedures, 277–278
- Stegostomatidae (zebra sharks), 47, 48
- Stingaree
  - lobed (*Urolophus lobatus*), 137, 156
  - masked (*Trygonoptera personata*), 153
- Stingarees (Urolophidae), 51–52
- Stingray
  - Atlantic (*Dasyatis sabina*), 137
  - bluntnose (*Dasyatis say*), 153
- Stingrays (Myliobatiformes), 51–52
- Stock assessment, 10–15, 205–240
  - age-structured models, 208
  - background and principles, 207–208
  - Bayesian approaches, 15
  - cohort models, Virtual Population Analysis, 220–224
  - conclusions and recommendations, 234–236
  - cross-comparison, 13–15
  - decision analysis approaches, 15
  - defined, 9, 10
  - delay-difference model (Deriso), 215–220
  - examples, 236
  - fishery dynamics, 11
  - maximum sustainable yield (MSY), 11–13
  - model complexity, 13–15
  - model fitting, 228–234
    - Bayesian estimation, 230–231
    - data quality, 231–233
    - linear regression, 229–230
    - time-series, 230
  - quantitative prediction, 10–11
  - surplus production (biomass dynamics) models, 208–212, see also Surplus production models

- uncertainty in, 11
- yield per recruit model, 212–215
- Stock collapse, 1–2
  - bycatch and, 2
  - mixed-species fishery mortality, 2
- Stock identification, 65
- Stock structure, molecular genetics, 79–96, see also Molecular markers
- Stratum boundaries, 272
- Stratum index, 271
- SUDAAN program, 280
- Surplus production models, 208–212
  - advantages and disadvantages, 212
  - data requirements, 211–212
  - dynamic, 207, 208
  - Fox model, 211
  - logistic growth/Schaefer model, 208–210
  - Pella–Tomlinson model, 211
  - principles of, 209
  - problems with, 207
- SURVAN program, 280
- Survey design, 271–273
- Survey index, 267–268
- Surveys
  - duration, 275–276
  - overall cost, 272
  - stratified random vs. systematic designs, 272
  - utility of, 267–269
- Survival, 72, see also Mortality
  - first-year, 194
  - to mean age at maturity, 194
  - respiratory mode and, 249
  - species comparisons, 249
  - stage-specific probabilities, 197–198
- SURVIV software, 178

Tagging, chemical, 116

Tag recovery studies

- in age/growth validation, 116
- approaches, 65
- assumptions, 73
- assumption violation in, 179
- background and principles, 59–60
- data collection and analysis, 63–73
- gear selectivity, 69–70
- growth rates, 66–69
- length/weight relationship, 66
- in mortality estimation, 176–180
- movement, 70–72
- problems of, 178
- relative abundance, 72
- sampling design, 64
- spatial and temporal distribution, 72
- species composition, 72–73
- survival/mortality, 72
- tag type and placement, 60–63, see also Tags

Tags

- archival, 74–75
  - pop-up, 74–75
- dart, 63
- internal anchor tag, 61–62, 63
  - body cavity type, 61–62
  - button type, 62
- Petersen disc tag, 61, 63
- Rototags, 62–63
  - Jumbo, 62–63
  - ORI, 62–63

Tail, family key, 42

Taxonomy

- classification, 23–24
- dissection, 32
- family key, 32–45

field identification, 28–30  
glossary/terminology, 24–28  
laboratory identification, 30  
orders and families, 45–53  
photography, 30  
specimen collection, preservation, and cataloguing, 31

Telemetry, 65, 180

Telephone sampling, 260–262

Terminal F assumption, 224

Terminology, taxonomic, 24–28

Territorial Use Rights in Fishing (TURFs), 300–301

Testis, sperm stages, 149

Tetracycline validation method, 117–118

Thornback ray (*Raja clavata*), 74, 136, 154

Thornback rays (Platyrrhinidae), 52

Thorns, caudal, in age and growth studies, 101

Thorns or fine denticles, family key, 35–36

Threatened species, 315–316

Thresher shark  
    bigeye (*Alopias supercilops*), 136  
    pelagic (*Alopius pelagicus*), 136

Thresher sharks (Alopiidae), 48

*Thunnus maccoyii* (southern bluefin tuna), 74

Tiger shark (*Galeocerdo cuvier*), 66, 142, 249

Time restrictions, 307–308

Time-series model fitting, 230

Tope shark (*Galeorhinus galeus*), 66, 153, 154, 155, 156

Torpedinidae (torpedo rays), 50

Torpediniformes (electric rays), 50  
    reproductive strategy, 136

Torpedo rays (Torpedinidae), 50

Total allowable catch (TAC), 304

Transmitted vs. reflected light, 105–106

Transverse vs. sagittal sectioning, 100–101

Trawl fishing gear, 247–248, 273, 289

*Triaenodon obesus* (whitetip reef shark), 146

Triakidae (houndsharks), 49, 50

*Triakis semifasciata* (leopard shark), 180

- rebound potential, 194
- staining methods, 111

Trip limits, 304

Trophonemata, reproductive strategy, 137

Trunk, family key, 39

*Trygonoptera personata* (masked stingaree)reproductive strategy, 153

Turtle Excluder Devices (TEDs), 247

  

Uncertainty, in stock assessment, 11

UN Fish Stocks Agreement, 299

United Nations Food and Agricultural Organization (FAO), 3, 298, 300

United States, reporting classes, 252

Urea content, 326

Urolophidae (stingarees), 51–52

*Urolophus lobatus* (lobed stingaree), 137, 153, 156

U.S. National Marine Fisheries Service, see NMFS entries

Use rights, 289–305

- limited entry, 301–302
- quantitative input (effort), 302–304
- quantitative output (catch quotas), 304–305
- territorial, 300–301

Utilization, 323–336

- consumptive, 325–331
  - cartilage, 329–330
  - fins, 326–328
  - liver, 330–331
  - meat, 325–326
  - peripheral products, 331
  - skin, 328
- non-consumptive, 332–333
  - catch-and-release fishing, 332–333
  - recreational diving, 332

## Validation

- absolute age vs. periodicity of ring formation, 116
- of age/growth findings, 115–119
- determinate methods (validation), 117–119
- indeterminate methods (verification), 116–117

Vernacular names vs. Latin binomial, 252

Vertebrae, transverse vs. sagittal sectioning, 100–101

Vertebral counts, 29–30

Vertebral growth, 99

Virtual Population Analysis, 180, 220–224

- example, 222–224

Vitamin A, liver as source, 330

## Viviparity

- aplacental, 136
- placental, 137–138

Von Bertalanffy growth function, 66–67, 119, 120–121, 168, 213–214

- sample size and, 121

Weasel sharks (Hemigaleidae), 49

## Web resources

- for age/growth studies, 123
- reproductive biology, 158

Wedgefishes (Rhincobatiformes), 51

Weight recording, 255–257

Weir & Cockerhan  $q$  estimator, 82

Whale shark (*Rhincodon typus*), 47, 309

Whiskery shark (*Furgaleus macki*), 152, 156

White shark (*Carcharodon carcharias*), 87, 90

Whitetip reef shark (*Triaenodon obesus*), 146

Wobbegongs (Orectolobidae), 47–48

World Conservation Union Shark Specialist Group, 2

Wright's  $F_{ST}$  statistic, 79

Yield per recruit model, 212–215

- advantages and disadvantages, 214–215
- data requirements and assumptions, 213

examples, 214

method, 213–214

Yolk sac, aplacental, 138

Young-of-the-year (YOY) tagging, 65

Zanobatoidei (stingrays), 51–52

Zebra sharks (Stegostomatidae), 47, 48